

Relativistic Modified Newtonian Dynamics from String Theory?

Nick E. Mavromatos and Mairi Sakellariadou

King's College London, Department of Physics, Strand WC2R 2LS, London, U.K.

We argue that TeVeS-like vector fields appear naturally in certain string theory backgrounds involving D0-branes, as a result of the recoil velocity field, expressing the interaction of neutrino string matter with point-like branes. However, the similarity with TeVeS models is restricted only to the bi-metric properties of space time, namely the difference of the background metric from the one felt by (some) matter fields interacting, in a topologically non-trivial manner, with the D0-brane defects. In our approach, neutrinos appear as dark matter candidates that could be “captured” by the D0 branes, as a result of stringy properties, and thus couple with the recoil-vector fields. Moreover, we argue in support of a possibly preferential rôle of neutrinos in inducing novel non-perturbative contributions to “vacuum” (dark) energy, in addition to their ordinary dark matter contribution. In fact, the rôle of neutrinos as providing substantial contributions to dark matter and dark energy components of the Universe, suggested by our approach, appears also to be necessitated by the need to reproduce the peaks in the Cosmic Microwave Background radiation spectrum, as claimed recently in the literature. Thus, our framework may be viewed as providing a microscopic explanation of such phenomenological conclusions concerning TeVeS-like, Lorentz-violating models.

PACS numbers: 98.80.-k, 95.36.+x, 95.35.+d, 95.85.Ry

Overwhelming experimental evidence from diverse sources [1], appear to provide convincing arguments in support of the existence of 70% of the universe’s total density in the form of dark energy, the fuel that drives acceleration, and 23% in an unknown form of weakly interacting Dark Matter (DM). However, the above studies are not model independent, as they are based on conventional Einstein/Friedmann-Lemaître-Robertson-Walker (FLRW) cosmologies. Deviations from these assumptions may lead to a different picture concerning the various contributions to the current-epoch energy budget of the universe. For instance, in non-equilibrium string theory models [2], it is possible to fit the data with the conventional Λ CDM model [3]. The important feature of these models is that the contributions to the universe’s energy budget, are coming from both dark energy and (dark and ordinary) matter terms.

More drastic conclusions can be drawn on the nature of the “dark sector” of the universe if one is prepared to abandon the Einstein gravity theory at galactic scales, in favour of a MODified Newtonian Dynamics (MOND) [4] below some universal gravitational acceleration scale a_0 . In such a case, the entire concept of DM may not be needed. Indeed, MOND theory, without DM at galactic scales, appears capable of explaining the rotational curves of galaxies, whose observed departure from the expected Keplerian law, in case all matter were luminous, led initially to the DM postulate. Although MOND originally appeared as a purely empirical (and not entirely covariant) theory, nevertheless recently Bekenstein [5] proposed a relativistic field theory version of it, involving TENSOR, VECTOR and SCALAR fields (TeVeS) of gravitational origin, reproducing the dynamics of MOND.

The TeVeS theory contains two metrics: $g_{\mu\nu}^E$, the “Einstein-frame” metric, which satisfies the canonically normalised Einstein-Hilbert action, and $g_{\mu\nu}$, that matter

“feels”; all geodesics are calculated in terms of $g_{\mu\nu}$. It is related to the Einstein-frame metric by [5]:

$$g_{\mu\nu} = e^{-2\Phi} g_{\mu\nu}^E + (e^{-2\Phi} - e^{2\Phi}) A_\mu A_\nu ; \quad (1)$$

A_μ (Φ) is the vector (scalar) field of the TeVeS theory. The presence of a vector field, A_μ , is crucial for consistency of the TeVeS models. Isotropy and homogeneity of the universe are not disturbed if A_μ has only a time-like component, which depends only on the cosmic time of the co-moving frame τ , namely $A_\mu = (A_0(\tau), 0, 0, 0)$, while internal consistency of the TeVeS theories implies a constraint on the fields magnitude [5]:

$$A_\mu A^\mu = -1 , \quad (2)$$

where the indices are contracted with respect the Einstein-frame metric. This constraint, of course, leads to Lorentz violation in the sense of its direction defining a “preferred frame”, thereby constituting a modern version of (isotropic) “aether” [6]. In a cosmological framework, with a scale factor $a(\tau)$, and Einstein metric such that $g_{00}^E = -a^{-2}(\tau)e^{2\Phi(\tau)}$, the constraint Eq. (2) implies that $A_0(\tau) = a(\tau)e^{\Phi(\tau)}$. The action of A_μ is described by a Maxwell-like kinetic term, in terms of its field strength ($F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$) plus a Lagrange multiplier term implementing the constraint, Eq. (2). The coefficient of the Maxwell term in the action is a free parameter.

In TeVeS the dynamics of Φ is described by an action including a non minimal coupling of the kinetic terms of Φ with another, non dynamical, scalar field μ , with a potential $V(\mu)$ whose form determines the phenomenology as far as the reproduction of the observed rotational curves of galaxies is concerned [5]. In Ref. [7] it was argued that TeVeS can reproduce the observed galactic power spectrum of baryonic fluctuations. This was a prediction of the DM hypothesis: just a baryonic universe predicts pronounced wiggles in the power spectrum

$\Delta \equiv k^3 P(k)/2\pi^2$, which are not observed. However, Ref. [7] argued that TeVeS could also explain the absence of the wiggles and reproduce the observed galaxy spectra without DM. It is the vector field of TeVeS that plays a crucial rôle in reproducing the observed spectra [8].

The arguments on the rôle of the vector field in reproducing large scale structures and the correct phenomenology seem to be generic, and not depending on the detailed Lagrangian of the original TeVeS theory. Only the basic important features of the theory, namely the existence of the two metrics and of the (Lorentz-violating) “aether-like” vector field, appear to be important in this respect. TeVeS can also reproduce [7] the observed positions of acoustic peaks in the CMB spectrum, only if a contribution from *massive neutrinos* is included, with a current density $\Omega_\nu \sim 0.15 - 0.17$, and masses ~ 2 eV, together with a *cosmological constant* contribution $\Omega_\Lambda \sim 0.80 - 0.78$, respectively, and ordinary (primarily baryonic) matter $\Omega_b \sim 5\%$. However, TeVeS theories have some drawbacks. They appear as “phenomenological”, not derived from an underlying microscopic theory, and they fail to explain the behaviour observed in certain systems, such as the bullet cluster [9], or galaxies claimed to be dominated by “dark matter” [10].

All the above features call for a microscopic explanation. Our aim is to argue in favour of a link between them, in the framework of some specific backgrounds of string theory [11]. The model involves two stacks of parallel eight-dimensional brane worlds embedded in a nine-dimensional bulk space. The bulk space-time is restricted in a specific way by two orientifolds, whilst the brane worlds can be compactified to three (spatial) dimensions in a specific way, whose details are not important here. The bulk involves D0-branes (D-particles), which can propagate in *both* the bulk and the brane worlds. They are viewed as point-like *defects* on the space-time. The D-particles have mass $m_D = M_s/g_s$ (M_s the string scale and $g_s < 1$ the string coupling). If M_s is at a TeV scale, then the mass of the defect is heavier. There are models in which $M_s = M_{\text{Pl}}$ (the four-dimensional Planck scale), in which case the D-particles have trans-Planckian mass.

One of the (compactified) three-branes plays the rôle of our observable universe, on which Standard Model matter lives. The latter is described by open strings ending to the D-brane world. Closed strings describe excitations in the gravitational multiplet of strings, and propagate in *both* the bulk and the brane. When the branes and the D-particles are *at rest* the vacuum is supersymmetric, with *zero vacuum energy*. Motion of branes breaks the target-space supersymmetry resulting in non-trivial contribution to the vacuum (“dark”) energy. Interaction of open strings on the brane with D-particles can be described by logarithmic field theory of ‘recoil’/impulse deformations of the pertinent σ -model describing stringy excitations on the D-particle defect [12]. The interaction is not a smooth scattering event. It involves the

capture of a string by the D-particle, which formally imply the change of world-sheet boundary conditions from Neumann to Dirichlet. Electric charge conservation implies that charged string excitations cannot be captured by the D-particle, due to the electromagnetic gauge symmetry $U_{\text{em}}(1)$. Since the D-particles also carry another $U(1)$ symmetry, which is unrelated to electromagnetism, flux conservation for that symmetry as well implies that an isolated D-particle, with a string emanating from it with one free end in the bulk space, *cannot* exist. Thus, from the elementary particles of the Standard Model, only electrically neutral particles can interact with D-particles.

As we shall discuss later on, *neutrinos* appear to have a preferential rôle in inducing novel non-perturbative contributions to the vacuum energy [13, 14], as compared to the other electrically neutral excitations that could in principle interact with D-particles. During each capture process, the massive D-particle defect recoils, and thus disturbs the surrounding space-time, resulting in metric distortions $\delta g_{\mu\nu}$. Such distortions are found by following conformal field theory methods on the world-sheet of the pertinent σ -model [12]. The vertex operator on the world-sheet boundary $\partial\Sigma$ reads

$$\int_{\partial\Sigma} u_i X^0 \Theta_\epsilon(X^0) \partial_n X^i ,$$

where ∂_n denotes normal world-sheet derivative, X^0 (X^i) obeys Neumann (Dirichlet) boundary conditions on the world-sheet, $u_i = \gamma v_i = \gamma g_s \Delta k_i / M_s$ is the “recoil” three-velocity of the D-particle defect during the capture process, $\gamma = (1 - \vec{v}^2)^{-1/2}$ is the Lorentz factor, and Δk_i is the momentum transfer of the stringy-matter excitation. The operator $\Theta_\epsilon(X^0)$, $\epsilon \rightarrow 0^+$, is a regularised Heaviside function, denoting the moment of impact, $X^0 = 0$ for definiteness [12]. World-sheet conformal invariance requires that a target-space metric deformation corresponds to a conformal-dimension-two operator of the σ -model. To find this deformation we first rewrite [12] the boundary deformation as a total world-sheet derivative in a target-space covariant form

$$\int_{\Sigma} \partial_{\hat{\alpha}} (u_\mu u_\nu X^\nu \Theta_\epsilon(u_\rho X^\rho) \partial^{\hat{\alpha}} X^\mu) ,$$

where $\hat{\alpha} = 1, 2$ is a world-sheet index, u_μ is a four-velocity,

$$u_\mu u^\mu = -1 . \quad (3)$$

For a linear dilaton $\Phi = -u_\mu X^\mu$ and σ -model frame Minkowski metric backgrounds [15], $e^{-\Phi}$ acts as the “Liouville dressing operator” [16], restoring conformal invariance, *i.e.*

$$\int_{\Sigma} e^{-\Phi} \partial_{\hat{\alpha}} (u_\mu u_\nu X^\nu \Theta_\epsilon(u_\rho X^\rho) \partial^{\hat{\alpha}} X^\mu)$$

has dimension two in the limit $\epsilon \rightarrow 0^+$:

$$V_{\text{bulk rec}} = \int_{\Sigma} e^{u_{\mu} X^{\mu}} u_{\mu} u_{\nu} \partial X^{\mu} \partial X^{\nu} \Theta_{\epsilon}(X^0) + \dots, \quad (4)$$

where the $\dots$ denote terms which vanish either upon using the world-sheet equations of motion, or upon taking the limit $\epsilon \rightarrow 0^+$. Indeed, due to the existence of a linear dilaton background, the operator $e^{-\Phi} = e^{u_{\mu} X^{\mu}}$ has conformal dimension $u_{\mu}(u^{\mu} - u^{\mu}) = 0$, thus it does not affect the overall conformal dimension two of the bulk recoil operator, Eq. (4), as a result of the $\partial X^{\mu} \partial X^{\nu}$ part.

The result of Eq. (4) points towards the existence of a target-space deformation due to the D-particle recoil:

$$\delta g_{\mu\nu} \propto e^{-\Phi} u_{\mu} u_{\nu} \Theta_{\epsilon \rightarrow 0}(u_{\rho} X^{\rho}).$$

However, to ensure a smooth connection with the flat metric at the origin of the boosted time $u_{\rho} X^{\rho} = 0$, it is necessary to impose the condition that $\delta g_{\mu\nu} = 0$ for $u_{\rho} X^{\rho} = 0$ (the reader should recall that at the co-moving frame of the recoiling defect $u_{\mu} X^{\mu} = X^0$, and thus the above condition is imposed at the origin of the co-moving frame target time). In the linear-dilaton scenario we employed above, this condition translates to the following metric, which describes the effects of the interaction of the stringy matter excitation with the D-particle defect on the surrounding space-time:

$$g_{\mu\nu}^{\text{matter}} = \eta_{\mu\nu} + (e^{\Phi} - e^{-\Phi}) u_{\mu} u_{\nu}, \quad \Phi = -u_{\mu} X^{\mu}.$$

The e^{Φ} correction corresponds to an operator on the world-sheet of the string with *zero* conformal dimension,

$$-u_{\mu}(-u^{\mu} - u^{\mu}) + 2 = 2(u_{\mu} u^{\mu} + 1) = 0,$$

due to Eq. (3). This latter deformation leads to departure of the associated σ -model from criticality.

To restore conformal invariance one can follow two approaches, which we shall only outline here, reserving detailed studies for a future publication. In view of the landscape scenario of string theory, both approaches could lead to acceptable in principle ground states of strings, which however violate Lorentz invariance, due to the preferred frame imposed by the recoil velocity field u_{μ} .

In the first approach, one stays within the Minkowski σ -model-frame metric background, and exploits the fact that the linear dilaton implies a sub-critical string with $Q^2 = u_{\mu} u^{\mu} = -1 < 0$. This can become conformal (critical) if one uses a *space-like* Liouville mode [16] ρ to “dress” the above-mentioned metric deformations by multiplication with exponential operators $e^{\alpha_i \rho}$, $i = 1, 2$, where α_i are the Liouville “anomalous” dimensions. In the presence of the world-sheet background charges in the (ρ, X^{μ}) extended target space time [16] of the Liouville-dressed world-sheet theory, ($|Q| = 1, u_{\mu}$), which induce world-sheet curvature terms of the form

$$\int_{\Sigma} d^2 \xi (u_{\mu} X^{\mu} + \rho) R^{(2)}$$

in the σ -model action. The conformal dimension of these operators is $\alpha_i(\alpha_i + 1)$ for each $i = 1, 2$. Restoration of conformal invariance requires that the total conformal dimension of the Liouville-dressed deformations is (1,1) in the (holomorphic, anti-holomorphic) world-sheet sectors. It is straightforward then to observe that the following dressed operators are conformal, amounting to the choices $\alpha_i = \pm|Q| = \pm 1$ in the respective Liouville anomalous dimensions,

$$V_{\text{dress}}^{\lambda\nu} u_{\lambda} u_{\nu} = \left(e^{-u_{\mu} X^{\mu} - \rho} - e^{u_{\mu} X^{\mu} + \rho} \right) \partial X^{\lambda} \bar{\partial} X^{\nu} u_{\lambda} u_{\nu}. \quad (5)$$

These imply a dressed target-space-time metric in the extended space-time (ρ, X) of the form:

$$\begin{aligned} g_{\mu\nu}^{\text{matter, dressed}}(\rho, X) &= \eta_{\mu\nu} + \left(e^{\Phi(\rho, X)} - e^{-\Phi(\rho, X)} \right) u_{\mu} u_{\nu}, \\ g_{\rho\mu} &= 0, \quad g_{\rho\rho} = +1, \quad \Phi(\rho, X) = -u_{\mu} X^{\mu} - \rho. \end{aligned} \quad (6)$$

The extra space-like Liouville mode may thus be given the physical interpretation of a bulk dimension, in which recoil of the defects *does not take place*. In this picture, our brane space time is located at, say, $\rho = 0$. This is only one example of a consistent conformal theory. In general, one may consider non-trivial σ -model metrics $G_{\mu\nu}$, in which case the associated dilatons will have a more complicated space time dependence. In this respect, one can discuss realistic FRW backgrounds, where however the dilaton potential (receiving contributions from higher string loops) plays a crucial rôle in the physics [2, 17]. We hope to come back to such issues in a forthcoming publication.

In the second approach to restoring conformal invariance, one remains within the original space time, but departs from the flat Minkowski background σ -model metric $\eta_{\mu\nu}$, by allowing a generic background $G_{\mu\nu}$. The latter can then be determined by the requirement of the vanishing of the associated Weyl anomaly coefficients [18] $\tilde{\beta}_{\mu\nu}^{G, \Phi}$. This, in turn, implies [18] that the associated σ -model renormalisation-group β -function for the deformed graviton vertex operator should equal an (infinitesimal) target-space diffeomorphism variation of the associated background field, *i.e.*,

$$\beta^i = -\delta g^i, \quad g^i = (G^{\mu\nu}, \Phi).$$

For the combination of background fields appearing in the expression for the “metric” felt by the matter interacting with the D-particle in our model:

$$g_{\mu\nu}^{\text{matter}} \equiv G_{\mu\nu} + (e^{\Phi} - e^{-\Phi}) u_{\mu} u_{\nu}, \quad (7)$$

one has for the associated world-sheet β -functions:

$$\beta_{\mu\nu}^{\text{g, matter}} = \beta_{\mu\nu}^G + 2\beta^{\Phi} \cosh(\Phi) u_{\mu} u_{\nu}, \quad (8)$$

assuming marginal (*i.e.*, world-sheet-renormalisation-group-scale independent) u_μ [12] (see, however, discussion below, where more general u_μ may appear). Requiring conformal invariance implies that the left-hand-side of Eq. (8) must have the form of the appropriate target space diffeomorphism variation $\delta g_{\mu\nu}^{\text{matter}}$. From Eq. (8) it is evident that it suffices to consider conformal invariance conditions for the dilaton Φ and σ -model-frame graviton $G_{\mu\nu}$ space time backgrounds, in the presence of the recoil deformations, which as discussed in Ref. [12], and will be reviewed below, act as if they are target space dynamical vector fields coupled to the metric $G_{\mu\nu}$. Indeed, in such a case

$$\beta_{\mu\nu}^G = -\delta G_{\mu\nu} \quad , \quad \beta^\Phi = -\delta\Phi \quad ,$$

and from Eq. (8) it follows trivially that

$$\begin{aligned} \beta_{\mu\nu}^{\text{matter}} &= -\delta G_{\mu\nu} - 2\cosh(\Phi)u_\mu u_\nu \delta\Phi \\ &= -\delta(G_{\mu\nu} + 2\sinh(\Phi)u_\mu u_\nu) \\ &= -\delta g_{\mu\nu}^{\text{matter}} \quad , \end{aligned} \quad (9)$$

which guarantees the world-sheet conformal-invariance of the matter metric Eq. (7).

In this article we shall not discuss in detail the existence of solutions to these constraints and their cosmological relevance. We only mention that in string models of the type considered in Ref. [11], where a population of D0-branes exists, it is possible to consider an isotropic situation, in which the statistical average $\langle\langle\ldots\rangle\rangle$ over the D0-branes of the spatial component of the recoil velocity vanishes, $\langle\langle u_i \rangle\rangle = 0$, $i = 1, 2, 3, \dots$. Effectively, if one ignores fluctuations of order $\langle\langle u_i u^i \rangle\rangle$, this may be considered equivalent to assuming only a temporal non-trivial component for u_μ , $u_0 \neq 0$, in agreement with the standard spatially isotropic TeVeS cosmologies [7, 8]. The constraint $u_\mu u^\mu = -1$, then, would imply that u_0 is a constant, as assumed above, only in cases where the temporal component of the background metric is -1 (as happens, for instance, in standard FLRW cosmologies in the appropriate time frame). In general, however, one may consider other frames, or more general situations, for instance the conformal time ones of Refs. [7, 8], in which the temporal component of the σ -model metric will not be -1 . In such cases, u_μ are not constants, but depend on the dilaton, the scale factor *etc.*, and the conformal invariance conditions get more complicated. We postpone a detailed discussion of such a situation for a forthcoming publication.

In either of the above ways of restoring the conformal invariance of the σ -model, we note the existence of two metrics, which is reminiscent of TeVeS theory, Eq. (1). One, is the σ -model background metric, metric, $G_{\mu\nu}$, which is related to an Einstein-frame metric $g_{\mu\nu}^E$ via [15]: $g_{\mu\nu}^E = e^{-4\Phi/(d-2)}G_{\mu\nu}$ in d -dimensional space-time. The other, is the metric $g_{\mu\nu}^{\text{matter}}$ describing the distortion of

the space-time surrounding the D-particle defect, as a result of its interaction with stringy matter:

$$g_{\mu\nu}^{\text{matter}} = e^{4\Phi/(d-2)}g_{\mu\nu}^E + (e^\Phi - e^{-\Phi})u_\mu u_\nu \quad . \quad (10)$$

The reader is invited to compare Eqs. (1) and (10); there are differences in the coefficients of the scalar field in the various exponentials, but the basic qualitative bi-metric features, are common. The dynamical scalar field of the TeVeS theory is thus played by the dilaton field in this model. The latter can also be responsible for yielding quintessence-like dark energy contributions [2, 3, 17]. We observe that for $d = 6$, one encounters a precise analogy with TeVeS models, which implies a Φ -independent electromagnetic fine-structure constant [5]. The velocity field u_μ , which is subject to the constraint Eq. (3) is *not* directly related to the dynamical vector field A_μ of the TeVeS theory, subject to the constraint, Eq. (2).

There is a dynamical gauge background field in the D-particle recoil model, which upon T -duality (which exchanges Neumann and Dirichlet boundary conditions) is related to the recoil deformation [12] :

$$\mathcal{A}_\mu = u_\mu \Phi \Theta(\Phi) \quad , \quad \Phi = -u_\mu X^\mu \quad , \quad (11)$$

with Lagrangian of the Born-Infeld type:

$$\frac{1}{g_s} \sqrt{\det(\eta_{\mu\nu} + 2\pi\alpha' g_s^2 F_{\mu\nu})} \ni \alpha' g_s^3 F_{\mu\nu} F^{\mu\nu} \quad ; \quad (12)$$

$F_{\mu\nu}(\mathcal{A})$ is the (Abelian) field strength. This holds for flat backgrounds. In general, for non-trivial σ -model-frame metric backgrounds $\eta_{\mu\nu} \rightarrow G_{\mu\nu}$ in (12), and there are also overall dilaton exponential factors, leading to relaxation dark energy terms [2, 11, 17].

In a galactic region, where we expect a statistically significant population of D-particles, there is a distribution of recoil velocities, thus one has to average Eq. (11) over such three-velocity distributions. We make the physically plausible assumption that such distributions are *isotropic* in space [11], $\langle\langle v_i \rangle\rangle = 0$. In this sense, the so-averaged gauge field depends solely on the target time $X^0 > 0$, and is of the form of the cosmological TeVeS field. In view of Eq. (3), the gauge field obeys a *gauge fixing* condition $\mathcal{A}_\mu \mathcal{A}^\mu = -\Phi^2$, which plays the rôle of the constraint Eq. (2). The indices are raised and lowered by the Einstein-frame metric $g_{\mu\nu}^E$. In the cosmological context of the TeVeS model, one may assume for g_{00} the form $g_{00}^E = a(\tau)^2 e^{-2\Phi} \eta_{00}$, with a the scale factor of the universe. This implies that $u^\mu = dX^\mu/d\tau$, where X^μ are σ -model-frame coordinates, and τ is a D-particle co-moving frame (proper) time. The co-moving cosmic time is then defined in the Einstein frame of the string by [15]: $d\tau = a e^{-\Phi} dX^0$, *i.e.*, $u^0 = a^{-1} e^\Phi$. Thus, like the TeVeS vector field, the Born-Infeld gauge field, describing the interaction of stringy matter with D-particle space-time defects, has the form

$$\langle\langle \mathcal{A}_0 \rangle\rangle = a e^{-\Phi} \Phi \Theta(\Phi) \quad . \quad (13)$$

From Eq. (12) we observe that the rôle of the parameter K of the TeVeS theory (the coefficient of the Maxwell-like kinetic term for the vector field) is played here by the g_s^3 , which is < 1 , for weak string coupling. For low-values of K , there is enhanced growth in the density perturbations [8].

We next remark on the special rôle of neutrinos in inducing non-perturbative cosmological-constant-type contributions to the vacuum energy in this picture. Neutrinos, as being electrically neutral, could be of the type of string excitations that interact with D-particles, leading to the existence of Lorentz-violating vector fields, Eq. (11), and the associated cosmological instabilities. The important point to notice [19] is that “flavor” is not *necessarily conserved* in such interactions. After the capture by the D-particle defect, the emerging stringy matter excitation could have a different flavor than what it had initially. Thus, the D-particle populations in galaxies act as a “medium” inducing flavor oscillations, in analogy with the celebrated Mikheev-Smirnov-Wolfenstein (MSW) effect [20]. Following Refs. [14, 19] we represent the ground state of the space-time with D-particle defects with which neutrinos interact as a Fock-space “flavor” vacuum $|0\rangle_f$, introduced in Ref. [13] in order to discuss quantisation of field-theories with mixing. Since in galaxies there is abundance of electrons (stable), there is a significant contribution to the vacuum energy coming from oscillations $\nu_e \leftrightarrow \nu_\mu$, since it is the electron current that couples to the muon neutrino current in Standard Model interactions. Thus, the dominant contributions to the vacuum energy can be computed in a two-generation neutrino model [13, 14]. Considering a cosmological space-time, say of FLRW type, and computing the average of the neutrino stress tensor with respect to the flavor vacuum ${}_f\langle 0|T_{\mu\nu}|0\rangle_f$, one finds that, in contrast to the usual mass-eigenstate Lorentz-invariant-vacuum case, where this quantity vanishes, there is a non-trivial, non-perturbative contribution to the vacuum energy [13]:

$$\rho_{\text{vac}} = \frac{2}{\pi} \sin^2 \theta \int_0^{K_0} d^3 k (\omega_{k,1} + \omega_{k,2}) |V_{\vec{k}}|^2, \quad (14)$$

K_0 a momentum cutoff, determining the relevant low-energy degrees of freedom, and

$$|V_{\vec{k}}|^2 \sim (m_1 - m_2)^2 / (4|\vec{k}|^2), \quad (15)$$

for large momenta, the “flavor” condensate. A consistent, and physically relevant choice of the cutoff is [14]

$$K_0 \sim m_1 + m_2. \quad (16)$$

This choice is compatible with our D-particle model since it implies that only the infrared neutrino modes, with momenta less than the typical mass scales $m_1 + m_2$, feel mostly the “D-particle medium” effects (being slow, they

have more time to interact with them). Thus, [14, 19]

$$\rho_{\text{vac}} \sim \frac{2}{\pi} \sin^2 \theta (m_1 - m_2)^2 (m_1 + m_2)^2 \sim \frac{2}{\pi} \sin^2 \theta (\Delta m_{12}^2)^2.$$

For $\Delta m_{12}^2 \sim 7 \times 10^{-5} \text{ eV}^2$, and $\sin^2 \theta \sim .3$, which are the measured values from atmospheric neutrino experiments, this leads to a dark-energy contribution from neutrinos

$$\Omega_{\Lambda}^{\nu_{\text{mixing}}} \sim 0.24. \quad (17)$$

The reader should keep in mind that there are additional time-dependent dark-energy contributions coming from the dilaton quintessence field [2, 17], which are not fully understood, as they include string loop corrections.

It is claimed [7] that the correct position of the CMB peaks is obtained if neutrino masses are of order 2 eV, their “dark matter” contribution is $\Omega_\nu \sim 0.15 - 0.17$, and the total $\Omega_\Lambda \sim 0.80 - 0.78$, respectively, with 5% ordinary matter. These features can be accommodated, if necessary, in our scenario. The presence of massive neutrino dark matter in galactic centres, would also contribute to the modification of the rotational curves of galaxies, on equal footing with the vector field. A detailed phenomenology of our model is left for future study.

A last but important point is that the D-particle medium consists of *supersymmetric* D0-brane defects, which are such that there is no contribution to the vacuum energy if interactions (recoil “movements”) with stringy matter are ignored. The D-particles are a sort of BPS states, which experience a zero net-force between them. Thus, the vacuum stress tensor of these supersymmetric defects vanishes and one avoids large isocurvature perturbations, which was a problem of other non-supersymmetric defects. Isocurvature perturbations are induced by the presence of “seeds” (any non-uniformly distributed form of energy, which contributes only a small fraction to the total energy density of the universe and which interacts with the cosmic fluid only gravitationally). Such perturbations are generated continuously and evolve according to inhomogeneous linear perturbation equations. The randomness of the non-linear seed evolution, which sources the perturbations, can destroy the coherence fluctuations in the cosmic fluid [21]. Moreover, if the seeds have non-linear dynamics, then the distribution of anisotropies leads to non-Gaussian statistics. Incidentally, small (unobservable) deviations from Gaussian statistics can also appear if the initial state of the field, responsible for the origin of fluctuations, is in a non-vacuum initial state [22]. Isocurvature perturbations and strong deviations from Gaussian statistics plague topological defect [23], as well as seed models in the context of pre-big-bang cosmology [24]. Severe constraints have been imposed [25] to the contribution of isocurvature perturbations to the CMB temperature anisotropies. In our model, D-particles could also lead to non-Gaussian signatures, due to $\langle\langle u_i^2 \rangle\rangle \neq 0$, which remain undetectable.

In our case, however, the presence of non-trivial dilaton fields may imply in general strong isocurvature perturbations, whose suppression amounts to severe constraints on model building. We hope to come back to such issues in the future.

In this work we have attempted to present models from string theory, leading to Lorentz-violating isotropic vector fields, and thus to a TeVeS theory. We have argued that one such model involves supersymmetric D-particles as a gravitational ground state “medium” on which stringy matter propagates. The interactions of matter with the defect involve “capture” of the matter string by the D0-brane, which leads to a different metric felt by the matter string as compared with the space-time background metric. The difference involves deformations by the recoil velocity field. We have presented arguments supporting the consistency of this bi-metric theory with world-sheet conformal invariance, which therefore makes it an acceptable string ground state. We have also argued that neutrinos are among the matter species that can interact in the above topologically non-trivial way (“capture”) with the D0 brane defects, and we have discussed the appearance of novel, non-perturbative contributions to the dark energy of the Universe as a result of the neutrino-D-particle interactions in such a model.

A final remark we would like to make is that the interactions of stringy matter with D-particles break, of course, the target-space supersymmetry of the vacuum, but this breaking is rather a *supersymmetry* obstruction [11], in the sense of yielding a non-supersymmetric spectrum of excitations. The breaking will be of order of the recoil velocity fluctuations $\langle\langle u_i^2 \rangle\rangle$, and thus for realistic situations very small. Hence, any phenomenologically realistic supersymmetry breaking on the D-brane world should be obtained from other means.

This work is partially supported by the European Union through the Marie Curie Research and Training Network *UniverseNet* (MRTN-CT-2006-035863).

-
- [1] A. G. Riess *et al.*, *Astrophys. J.* **560**, 49 (2001); A. G. Riess *et al.*, [arXiv:astro-ph/0611572](#); D. N. Spergel *et al.*, [arXiv:astro-ph/0603449](#); D. J. Eisenstein *et al.*, *Astrophys. J.* **633** (2005) 560; S. Cole *et al.* *Mon. Not. R. Astron. Soc.* **362**, 505 (2005).
 - [2] G. A. Diamandis *et al.*, *Phys. Lett. B* **642**, 179 (2006).
 - [3] J. R. Ellis, N. E. Mavromatos, V. A. Mitsou and D. V. Nanopoulos, *Astropart. Phys.* **27**, 185 (2007).
 - [4] M. Milgrom, *Astrophys. J.*, **270**, 365 (1983).
 - [5] J. D. Bekenstein, *Phys. Rev. D* **70**, 083509 (2004); R. H. Sanders, *Astrophys. J.*, **473**, 117 (1996).
 - [6] T. Jacobson and D. Mattingly, *Phys. Rev. D* **70**, 024003 (2004).
 - [7] C. Skordis, D. F. Mota, P. G. Ferreira and C. Boehm, *Phys. Rev. Lett.* **96**, 011301 (2006).
 - [8] S. Dodelson and M. Liguori, *Phys. Rev. Lett.* **97**, 231301 (2006).
 - [9] D. Clowe, A. Gonzalez and M. Markevitch, *Astrophys. J.* **604**, 596 (2004).
 - [10] E. O’ Sullivan and T. J. Ponman, *Mon. Not. R. Astron. Soc.* **354**, 935 (2004); I. Ferreras, P. Saha and L. L. R. Williams, *Astrophys. J.* **623**, L5 (2005).
 - [11] J. R. Ellis, N. E. Mavromatos and M. Westmuckett, *Phys. Rev. D* **70**, 044036 (2004); J. R. Ellis, N. E. Mavromatos, D. V. Nanopoulos and M. Westmuckett, *Int. J. Mod. Phys. A* **21** (2006) 1379.
 - [12] I. I. Kogan, N. E. Mavromatos and J. F. Wheeler, *Phys. Lett. B* **387**, 483 (1996); N. E. Mavromatos and R. J. Szabo, *Phys. Rev. D* **59**, 104018 (1999); *JHEP* **0110**, 027 (2001).
 - [13] M. Blasone *et al.*, *Phys. Lett. A* **323**, 182 (2004); A. Capolupo, S. Capozziello and G. Vitiello, [arXiv:astro-ph/0602467](#).
 - [14] G. Barenboim and N. E. Mavromatos, *Phys. Rev. D* **70**, 093015 (2004).
 - [15] I. Antoniadis, C. Bachas, J. R. Ellis and D. V. Nanopoulos, *Nucl. Phys. B* **328**, 117 (1989).
 - [16] F. David, *Mod. Phys. Lett. A* **3**, 1651 (1988); J. Distler and H. Kawai, *Nucl. Phys. B* **321**, 509 (1989).
 - [17] M. Gasperini, F. Piazza and G. Veneziano, *Phys. Rev. D* **65**, 023508 (2002).
 - [18] R. R. Metsaev and A. A. Tseytlin, *Nucl. Phys. B* **293**, 385 (1987).
 - [19] N. E. Mavromatos and S. Sarkar, *Phys. Rev. D* **74**, 036007 (2006).
 - [20] L. Wolfenstein, *Phys. Rev. D* **17**, 2369 (1978); S. P. Mikheev and A. Y. Smirnov, *Sov. J. Nucl. Phys.* **42**, 913 (1985) [*Yad. Fiz.* **42**, 1441 (1985)].
 - [21] M. Sakellariadou, [astro-ph/9911497](#).
 - [22] J. Martin, A. Riazuelo and M. Sakellariadou, *Phys. Rev. D* **61** 083518 (2000); A. Gangui, J. Martin and M. Sakellariadou, *Phys. Rev. D* **66** 083502 (2002).
 - [23] R. Durrer and M. Sakellariadou, *Phys. Rev. D* **56** 4480 (1997); R. Durrer, M. Kunz, C. Lineweaver and M. Sakellariadou, *Phys. Rev. Lett.* **79** 5198 (1997).
 - [24] R. Durrer, M. Gasperini, M. Sakellariadou and G. Veneziano, *Phys. Rev. D* **59** 043511 (1999); *Phys. Lett. B* **436** 66 (1998).
 - [25] R. Bean, J. Dunkley and E. Pierpaoli, *Phys. Rev. D* **74**, 063503 (2006).